

COMPOUND SODIUM LACTATE (HARTMANN'S) AND GLUCOSE INTRAVENOUS INFUSION

Consumer Medicine Information

What is in this leaflet?

This leaflet answers some common questions about Compound Sodium Lactate (Hartmann's) and Glucose IV Infusion. It does not contain all of the available information.

All medicines have risks and benefits. Your doctor has prescribed Hartmann's and Glucose IV Infusion for you in the knowledge that the benefits outweigh any possible risks. It does not take the place of talking to your doctor, nurse or pharmacist. If you have any concerns about receiving this medicine, ask your health care professional.

What Hartmann's and Glucose IV Infusion is used for

Hartmann's and Glucose IV Infusion is used to replace body fluid and mineral salts that may be lost for a variety of medical reasons. Hartmann's and Glucose IV Infusion is especially suitable when the losses result in too much acid being present in the blood. Glucose is a source of energy and is often used when a patient is recovering from a surgical operation.

Before you are given Hartmann's and Glucose IV Infusion

Hartmann's and Glucose IV Infusion should not be given to you if:

- you have congestive heart failure (constant wheezing, shortness of breath)
 - you have ischaemic stroke (have had a stroke)
 - you have liver disease as a consequence of alcoholism, or
 - you have severe reduction in kidney function
 - you have already too much glucose in your blood, or are not passing urine
- Also, the product should not be used if the expiry date printed on the bottom of the bag has passed, or if the packaging is torn or shows any signs of tampering.

You must tell your doctor if you:

- have heart disease
- have any other illness
- are taking other medicines including anything that you bought without a prescription from your pharmacy, supermarket or health food shop
- are pregnant
- are breast feeding
- are diabetic
- you have, at any time, had an allergic reaction to Hartmann's IV Infusion or to corn or corn products

How Hartmann's and Glucose IV Infusion is given

How much will be given:

Your doctor will decide how much Hartmann's and Glucose IV infusion will be given to you, which depends on your needs and medical condition. The solution is delivered by intravenous infusion (drip); therefore it should be given by using special equipment and attended by a health professional (doctor, trained nurse).

How it is given:

Hartmann's and Glucose IV infusion will be given at a slow rate of injection (drip) by a health professional. Usually, you will need to stay in a health institution (hospital, clinic, nursing home, etc). Special medical equipment is used to deliver the medicine into your bloodstream. This delivery should be attended by a health professional. A cannula (administration needle) is placed in a vein by your doctor or nurse. The infusion is for single use, and for one person only.

Hartmann's and Glucose IV Infusion must be used once only. Any unused portion must be discarded.

In case of overdose

Overdose with Hartmann's and Glucose IV Infusion is rare, and as your doctor has had experience in the use of this medicine, it is unlikely that you will be given an overdose. However, in the unlikely event that overdose does occur, the infusion will be discontinued and other treatment may be needed. You may experience some of the effects listed under "Side Effects" below.

While you are receiving Hartmann's and Glucose IV Infusion

Frequent clinical evaluation and laboratory tests may be required. As Hartmann's and Glucose IV Infusion is normally given in a hospital, your nurse or carer will make appropriate records during your treatment and will note any unexpected effects you may experience.

Side effects

As with any medicine, some side effects may occur. Some swelling of the hands, ankles and feet may be experienced due to retention fluid in your body. In rare circumstances, this may also involve the lungs, which may cause some breathing difficulty. Other symptoms may include nausea, vomiting, headache, dizziness, drowsiness or confusion. Inflammation at the site of injection, or swelling of your veins near the site of injection is also possible. Always tell your doctor or nurse immediately if you have any unexpected effects during or after receiving Hartmann's and Glucose IV Infusion.

Product descriptions

What Hartmann's and Glucose IV Infusion looks like

It is a clear colourless, or slightly yellow, solution packaged in Viaflex plastic bags. It is a sterile solution and intended for slow intravenous injection.

What is in Hartmann's and glucose IV Infusion?

The active components of Hartmann's and Glucose IV Infusion are: sodium lactate, potassium chloride, sodium chloride, calcium chloride and glucose. It is formulated in Water for Injections

How to store Hartmann's and Glucose IV Infusion

Hartmann's and Glucose IV Infusion should be stored below 30°C. Do not freeze. Do not expose to heat. Keep out of reach of children.

Where can you get more information?

You can get more information from your doctor or pharmacist.

Name and address of the manufacturer (sponsor)

Baxter Healthcare Pty. Ltd.,
1 Baxter Drive,
Old Toongabbie N.S.W., 2146 Australia
www.baxterhealthcare.com.au.

This Consumer Medication Information was prepared in February 2002 and revised on 23rd December, 2002

88-19-01-096

Baxter and Viaflex are registered trademarks of Baxter International, Inc.